



The TIBET AS+MD Project; progress report 2011

M. AMENOMORI¹, X. J. BI², D. CHEN³, W. Y. CHEN², S. W. CUI⁴, DANZENGLUOBU⁵, L. K. DING², X. H. DING⁵, C. F. FENG⁶, ZHAOYANG FENG², Z. Y. FENG⁷, Q. B. GOU², H. W. GUO⁵, Y. Q. GUO², H. H. HE², Z. T. HE^{4,2}, K. HIBINO⁸, N. HOTTA⁹, HAIBING HU⁵, H. B. HU², J. HUANG², W. J. LI^{2,7}, H. Y. JIA⁷, L. JIANG², F. KAJINO¹⁰, K. KASAHARA¹¹, Y. KATAYOSE¹², C. KATO¹³, K. KAWATA³, LABACIREN⁵, G. M. LE², A. F. LI^{14,6,2}, C. LIU², J. S. LIU², H. LU², X. R. MENG⁵, K. MIZUTANI^{11,15}, K. MUNAKATA¹³, H. NANJO¹, M. NISHIZAWA¹⁶, M. OHNISHI³, I. OHTA¹⁷, S. OZAWA¹¹, X. L. QIAN^{6,2}, X. B. QU², T. SAITO¹⁸, T. Y. SAITO¹⁹, M. SAKATA¹⁰, T. K. SAKO¹², J. SHAO^{2,6}, M. SHIBATA¹², A. SHIOMI²⁰, T. SHIRAI⁸, H. SUGIMOTO²¹, M. TAKITA³, Y. H. TAN², N. TATEYAMA⁸, S. TORII¹¹, H. TSUCHIYA²², S. UDO⁸, H. WANG², H. R. WU², L. XUE⁶, Y. YAMAMOTO¹⁰, Z. YANG², S. YASUE²³, A. F. YUAN⁵, T. YUDA³, L. M. ZHAI², H. M. ZHANG², J. L. ZHANG², X. Y. ZHANG⁶, Y. ZHANG², YI ZHANG², YING ZHANG², ZHAXISANGZHU⁵, X. X. ZHOU⁷ (THE TIBET AS γ COLLABORATION)

¹Department of Physics, Hirosaki University, Hirosaki 036-8561, Japan

²Key Laboratory of Particle Astrophysics, Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, China

³Institute for Cosmic Ray Research, University of Tokyo, Kashiwa 277-8582, Japan

⁴Department of Physics, Hebei Normal University, Shijiazhuang 050016, China

⁵Department of Mathematics and Physics, Tibet University, Lhasa 850000, China

⁶Department of Physics, Shandong University, Jinan 250100, China

⁷Institute of Modern Physics, SouthWest Jiaotong University, Chengdu 610031, China

⁸Faculty of Engineering, Kanagawa University, Yokohama 221-8686, Japan

⁹Faculty of Education, Utsunomiya University, Utsunomiya 321-8505, Japan

¹⁰Department of Physics, Konan University, Kobe 658-8501, Japan

¹¹Research Institute for Science and Engineering, Waseda University, Tokyo 169-8555, Japan

¹²Faculty of Engineering, Yokohama National University, Yokohama 240-8501, Japan

¹³Department of Physics, Shinshu University, Matsumoto 390-8621, Japan

¹⁴School of Information Science and Engineering, Shandong Agriculture University, Taian 271018, China

¹⁵Saitama University, Saitama 338-8570, Japan

¹⁶National Institute of Informatics, Tokyo 101-8430, Japan

¹⁷Sakushin Gakuin University, Utsunomiya 321-3295, Japan

¹⁸Tokyo Metropolitan College of Industrial Technology, Tokyo 116-8523, Japan

¹⁹Max-Planck-Institut für Physik, München D-80805, Deutschland

²⁰College of Industrial Technology, Nihon University, Narashino 275-8576, Japan

²¹Shonan Institute of Technology, Fujisawa 251-8511, Japan

²²RIKEN, Wako 351-0198, Japan

²³School of General Education, Shinshu University, Matsumoto 390-8621, Japan

takita@icrr.u-tokyo.ac.jp

DOI: 10.7529/ICRC2011/V06/0351

Abstract: We plan to build a large (approximately 10,000 m²) water-Cherenkov-type muon detector array under the existing Tibet air shower array at 4,300 m above sea level, to observe 10-1000 TeV gamma rays from cosmic-ray accelerators in our Galaxy with wide field of view at very low background level. The gamma-ray induced air shower has significantly less muons compared with a cosmic-ray induced one. Therefore, we can effectively discriminate between primary gamma rays and cosmic-ray background events by means of counting the number of muons in an air shower by the muon detector array. We will make a progress report on the project, as some part of it is under construction now.

Keywords: Tibet, cosmic rays, gamma rays, muon, water Cherenkov

1 Introduction

The HESS group with 4 large IACTs in Namibia, reported on discovery of new 14 gamma-ray sources by galactic plane survey [1] in 2005. Surprisingly, most of them are UNIDentified (UNID) sources and faint in X-rays or other wavelengths. As the HESS survey was limited within the galactic plane in the southern hemisphere due to its narrow field of view, the importance of a wide field-of-view unbiased survey is emphasized. Furthermore, many of the 14 sources have a harder energy spectrum (indices: -1.8 to -2.8) at TeV energies than the standard candle Crab (index: -2.6). The energy spectra turned out to extend up to approximately 10 TeV. Cosmic rays are supposed to be accelerated up to the knee energy region at supernova remnants (SNRs) in our galaxy. Therefore, we naturally expect gamma rays in the 100 TeV region (10-1000TeV) which originate in π^0 decays produced by the accelerated cosmic rays interacting with matter surrounding the SNRs. The gamma ray emission of electron origin might be highly suppressed in the 100 TeV region due to rapid decrease of inverse-Compton cross section by the Klein-Nishina effect as well as synchrotron radiation energy loss in the magnetic field around the SNRs. Multi-wavelength (radio, X-ray, gamma-ray) observations are essential to discriminate between the two processes.

2 Tibet Air Shower Experiment

The Tibet air shower array has been in operation at Yangba-jing (90°31' E, 30°06' N; 4300 m above sea level) in Tibet, China since 1990. It has continuously made a wide field-of-view (approximately 2 steradian) observation of cosmic rays and gamma rays in the northern sky.

The Tibet I array was constructed in 1990 [2], and it was gradually upgraded to the Tibet II by 1994 which consisted of 185 fast-timing (FT) scintillation counters placed on a 15 m square grid covering 36,900 m², and 36 density (D) counters around the FT-counter array. Each counter has a plastic scintillator plate of 0.5 m² in area and 3 cm in thickness. All the FT counters are equipped with a fast-timing 2-inch-diameter photomultiplier tube (FT-PMT), and 52 out of 185 FT counters are also equipped with a wide dynamic range 1.5-inch-diameter PMT (D-PMT) by which we measure up to 500 particles which saturates FT-PMT output, and all the D-counters have a D-PMT. A 0.5 cm thick lead plate is put on the top of each counter in order to increase the counter sensitivity by converting gamma rays into electron-positron pairs in an electromagnetic shower. The mode energy of the triggered events in Tibet II is ~ 10 TeV/ ~ 7 TeV for cosmic rays/gamma rays [3, 4].

In 1996, we added 77 FT counters with a 7.5 m lattice interval to a 5,200 m² area inside the northern part of the Tibet II array. We called this high-density array Tibet HD [5]. The mode energy of the triggered events in Tibet HD is ~ 3 TeV/ ~ 2 TeV for cosmic rays/gamma rays. In the late fall

of 1999, the array was further upgraded by adding 235 FT-counters to increase the high-density area from 5,200 m² to 22,050 m². We call this array and further upgraded one Tibet III [6]. In 2002, all of the 36,900 m² area was

covered by the high-density array by adding 200 FT-counters. Finally we set up 56 FT-counters around the 36,900 m² high density array and equipped 8 D-counters with FT-PMT in 2003. At present, the Tibet air shower array consists of 761 FT-counters (249 of which have a D-PMT) and 28 D-counters [7].

The performance of the Tibet air shower array has been well examined by observing the Moon's shadow (approximately 0.5 degree arc in diameter) in cosmic rays [6, 8]. The deficit map of cosmic rays around the Moon demonstrates the angular resolution to be around 0.9° at a few TeV for the Tibet III array. The pointing error is estimated to be less than $\sim 0.01^\circ$ by displacement of the shadow's center from the apparent center in the north-south direction, as the east-west component of the geomagnetic field is very small at the experimental site. On the other hand, the shadow center displacement in the east-west direction due to the geomagnetic field enables us to spectroscopically estimate the energy scale uncertainty less than $\pm 12\%$ [8].

The first celestial TeV gamma-ray signal from the Crab (the standard candle in gamma-ray astronomy) was detected by an imaging air Cherenkov telescope (IACT) developed by the Whipple group in 1989 [9]. The pioneering work opened a new energy window in astronomy.

Ten years later in 1999, we succeeded in observing multi-TeV gamma-ray signal from the Crab Nebula at 5.5σ confidence level, using the Tibet HD array [5]. This was the first detection of multi-TeV gamma-ray signal by a conventional air shower array. Subsequently, in 2000, we reported on detection of multi-TeV gamma rays successfully at 3.7σ level from Mrk501 which was in a highly flaring state during March 1997 and August 1997 [3]. We also succeeded in observing multi-TeV gamma-ray flares at 5.1σ level from Markarian 421 which was in a very active phase during the year 2000 and 2001 [6]. Multi-TeV gamma-ray signal was successfully detected from the Crab by the Tibet-III array as well [8].

We searched for TeV steady point sources in the northern sky. No statistically significant point source is found except for well established Crab and Mrk421 [10].

Search for steady PeV gamma-ray emission from the Monogem ring region is done with the Tibet data taken from 1997 to 2004 [11]. No evidence for statistically significant gamma-ray signal is found in the Monogem ring where the MAKET-ANI experiment recently claimed a positive detection of PeV high-energy cosmic radiation [12], although our flux sensitivity is approximately 10 times better than MAKET-ANI's.

The 2-dimensional anisotropy ($\sim \pm 0.2\%$) in the equatorial coordinates are obtained for the first time in the multi-TeV energy range [13]. Furthermore, we discovered a new diffuse cosmic-ray anisotropy (approximately 0.2 %

level excess) in the Cygnus region [14], although we cannot judge currently whether it is caused by gamma rays and/or a local cosmic-ray anisotropy. Overlapping the large-scale anisotropy in the Cygnus region, a few spacially separated enhancements of smaller scale were observed. These small-scale excesses ($\sim 2^\circ$) hint at the extended gamma-ray emission[14]. The two of the small-scale excesses were shortly established[15], [16] as extended gamma-ray sources by the Milagro experiment capable of hadron/gamma-ray discrimination. Recently, we found a positive statistical correlation between the Fermi bright Galactic sources and our TeV gamma-ray excesses[17].

We also searched for multi-TeV diffuse gamma rays from the galactic plane [4, 18]. As there was no significant signal, flux upper limits were obtained.

Meanwhile, the Milagro group recently claimed detection of TeV diffuse gamma-ray signal in the Cygnus region along the galactic plane [19]. Currently, we can not draw any clear conclusion on the Milagro result, as the Tibet air shower array is unable to distinguish between cosmic gamma rays and hadrons in the multi-TeV energy region at present.

To positively observe gamma rays in the 100 TeV region with much better sensitivity than Tibet III, we plan to add a muon detector array to the air shower array. Gamma-ray induced electromagnetic air showers are muon-poor, while cosmic-ray induced hadronic ones are accompanied by many muons. This enables us to separate gamma rays from cosmic rays. Our current plan [20, 21] relevant to gamma-ray astronomy above 10 TeV is Tibet AS (Air Shower array with $83,000\text{m}^2$ in area) + MD (Muon Detector array with $\sim 10,000\text{m}^2$ in area under Tibet AS). Each muon detector is a waterproof concrete pool, 7.2 m wide \times 7.2 m long \times 1.5 m deep in size, equipped with two 20 inch-diameter photomultiplier tubes (PMTs), i.e., HAMAMATSU R3600. The Tibet MD array are made up of 192 muon detectors set up 2.5 m underground. Its total effective area amounts approximately to $10,000\text{m}^2$ for muons with energies more than $\sim 1\text{GeV}$. Our current MC simulation predicts that the cosmic-ray background events will be rejected by approximately 99.99% at 100 TeV using full-scale ($10,000\text{m}^2$) MD array. the full-scale MD array will improve the sensitivity to gamma-ray sources by more than an order of magnitude.

To confirm the hadron rejection power with this full-scale MD array, we constructed a prototype muon detector in 2007.

3 Prototype Muon Detector

In the late fall of 2007, we constructed a prototype water Cherenkov muon detector (approximately 100m^2) at $\sim 90\text{m}$ away from the center of the existing Tibet airshower array.

The muon detector is made from reinforced concrete and composed of two water pool cells located at 2.5 m under




Figure 1: From [21]. Charge distribution recorded by the sum of two PMTs in a cell of the prototype muon detector. A peak of charge distribution is defined as one muon.

the ground. Each cell is filled up with water of 1.5 m in depth, $7.2\text{m} \times 7.2\text{m}$ in area, equipped with three $20''\phi$ downward facing PMTs (HAMAMATSU R3600). Among the 3 PMTs, one is covered by a black sheet with $\sim 1\%$ light transmission to effectively reduce the PMT gain (wider dynamic range), although we do not use it in this paper. The timing and charge information for each PMT is recorded by a trigger generated from surface scintillation counter array. We started the data taking in December, 2007.

4 Results and Discussions

Figure 1 shows the charge distribution by the sum of two PMTs in a cell (52m^2) extracted from air shower data. In this figure, a clear peak around 28 pC indicates the average charge when one particle pass through the muon detector cell. We calculate the muon number assuming this peak is defined as one muon.

Figure 2 and Figure 3 show the muon number distribution recorded by the prototype muon detector in an air shower at 10 TeV region ($100 < \Sigma\rho < 300$) and 100 TeV region ($1000 < \Sigma\rho < 3000$), respectively, where $\Sigma\rho$ means the sum of the number of particles per m^2 for each scintillation detector (10 TeV corresponds to $\Sigma\rho \sim 100$ roughly). As for the air shower MC simulation, air shower events are generated using CORSIKA code with QGSJET01c model for hadronic interactions assuming the cosmic-ray composition estimated mainly from direct observational data. Our MC simulation reproduces well the experimental data obtained by the prototype muon detector. Following the success, construction of some part of the full-scale muon detectors started in 2010.

5 Summary

In 2007, we constructed a prototype water Cherenkov muon detector (approximately 100m^2) under the existing Tibet air shower array. We have been successfully taking




Figure 2: Number of muons recorded by the prototype muon detector in an air shower at 10 TeV region ($100 < \Sigma\rho < 300$). The filled diamonds indicate experimental data, while the solid histograms are the MC simulation. Air showers with null muons detected by the muon detector are plotted at number of muons of 0.1.




Figure 3: Number of muons recorded by the prototype muon detector in an air shower at 100 TeV region ($1000 < \Sigma\rho < 3000$). The filled diamonds indicate experimental data, while the solid histograms are the MC simulation. Air showers with null muons detected by the muon detector are plotted at number of muons of 0.1.

data since December in 2007. The preliminary analysis shows that the data is in good agreement with our MC simulation. We are ready for the full-scale ($10,000 \text{ m}^2$) underground water Cherenkov muon detector array, some part of which is under construction now.

Acknowledgments

The collaborative experiment of the Tibet Air Shower Arrays has been performed under the auspices of the Ministry of Science and Technology of China and the Ministry of Foreign Affairs of Japan. This work was supported in part by a Grant-in-Aid for Scientific Research on Priority Areas from the Ministry of Education, Culture, Sports, Science and Technology, by Grants-in-Aid for Science Research from the Japan Society for the Promotion of Science in Japan, and by the Grants from the National Natural Science Foundation of China and the Chinese Academy of Sciences.

References

- [1] Aharonian, F.A. et al., *Science* **307**, 1938 (2005).
- [2] Amenomori, M. et al., *Phys. Rev. Lett.* **532**, 2468 (1992).
- [3] Amenomori, M. et al., *Astrophys. J.* **532**, 302 (2000).
- [4] Amenomori, M. et al., *Astrophys. J.* **580**, 887 (2002).
- [5] Amenomori, M. et al., *Astrophys. J.* **525**, L93 (1999).
- [6] Amenomori, M. et al., *Astrophys. J.* **598**, 242 (2003).
- [7] Amenomori, M. et al., *Astrophys. J.* **678**, 1165 (2008).
- [8] Amenomori, M. et al., *Astrophys. J.* **692**, 61 (2009).
- [9] Weeeks, T.C. et al., *Astrophys. J.* **342**, 379 (1989).
- [10] Amenomori, M. et al., *Astrophys. J.* **633**, 1005 (2005).
- [11] Amenomori, M. et al., *Astrophys. J.* **635**, L53 (2005).
- [12] Chilingarian, A. et al., *Astrophys. J.* **597**, L129 (2003).
- [13] Amenomori, M. et al., *Astrophys. J.* **626**, L29 (2005).
- [14] Amenomori, M. et al., *Science* **314**, 439 (2006).
- [15] Abdo, A.A. et al., *Astrophys. J.* **658**, 776 (2007).
- [16] Abdo, A.A. et al., *Astrophys. J.* **664**, L91 (2007).
- [17] Amenomori, M. et al., *Astrophys. J.* **709**, L6 (2010).
- [18] Amenomori, M. et al., *Advances in Space Research* **37**, 1932 (2006).
- [19] Atkins, R. et al., *Phys. Rev. Lett.* **95**, 251103 (2005).
- [20] Amenomori, M. et al., *AIP Conf. Proc.* **1085**, 723 (2008).
- [21] Amenomori, M. et al., *Journal of Physics: Conference Series* **120**, 062024 (2008).